

# LEGIBILITY NOTICE

A major purpose of the Technical Information Center is to provide the broadest dissemination possible of information contained in DOE's Research and Development Reports to business, industry, the academic community, and federal, state and local governments.

Although a small portion of this report is not reproducible, it is being made available to expedite the availability of information on the research discussed herein.

TITLE    **OSCILLATIONS OF SOLAR MODELS WITH INTERNAL ELEMENT DIFFUSION**

AUTHOR(S)    **Arthur N. Cox, T-6  
Joyce A. Guzik, T-6, Iowa State Univ.  
Russell B. Kidman, A-4**

SUBMITTED TO    **Proceedings of Solar Radiative Oupput Variations  
Boulder, Colorado, Nov. 9-11, 1987**

**DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U. S. Government retains a non-exclusive, royalty-free license to publish or reproduce reprinted forms of this contribution, or to allow others to do so, for U. S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U. S. Department of Energy.

---

**MASTER**

**Los Alamos** Los Alamos National Laboratory  
Los Alamos, New Mexico 87545

# OSCILLATIONS OF SOLAR MODELS WITH INTERNAL ELEMENT DIFFUSION

Arthur N. Cox<sup>1</sup>, Joyce A. Guzik<sup>1,2</sup>, and Russell B. Kidman<sup>1</sup>

<sup>1</sup>Los Alamos National Laboratory

<sup>2</sup>Astronomy Program, Department of Physics  
Iowa State University

Two precision solar models have been constructed with the Iben evolution program, one with no diffusion of the internal atomic nuclei, and another that includes the effects of gravitational settling, thermal diffusion, and concentration gradient diffusion on the element abundances. Our pressure and energy equation of state and opacity have been specially fit to the latest theoretical data. Then the opacity at the bottom of the convection zone was increased 15 percent (within its theoretical uncertainty) to allow a better agreement with the observed solar p-mode frequencies. The theoretical p-mode frequencies are still too small by about 10-20 microhertz. The solar mass, radius, luminosity, and age of  $4.6 \pm 0.1$  billion years were matched for both evolution runs so that the theoretical error of the predicted oscillation frequencies was less than one microhertz. Parameters that were adjusted to match with the  $M_{\odot}$ , the observed  $R_{\odot}$ , and  $L_{\odot}$  were the initial helium abundance and the ratio of the mixing length to the pressure scale height in the normal convection theory. The original helium mass fractions of the mixture were 0.300 and 0.298 for the no-diffusion and diffusion models, respectively. The diffusion model evolved to a surface  $Y=0.263$  at the solar age, and the original  $Z$  value of 0.0200 decreased to 0.0179 by diffusive settling. Our equation of state gives pressures that average a few percent too large throughout the sun, and therefore more helium, with its lower pressure per gram than hydrogen, is needed to compensate for this error. We estimate that the true helium mass fraction is about 0.02 smaller than what we use. The bottom of our convection zone is hotter than for most previous models because we have tried to match the p-mode frequencies better, but it is probably not hot enough to deplete the surface lithium abundance to the observed value unless there is overshooting below the bottom of the convection zone by about half a pressure scale height. It is possible that theoretical uncertainties in the pressure equation of state will always prevent matching the helium abundance ( $Y$ ) and the p-mode frequencies to better than 0.02 and 10 microhertz, respectively. A discussion is given of the agreement of asymptotic theory p-mode frequency separations and those directly calculated with nonadiabatic theory. Calculation of g-mode

solutions shows that they do not have equal spacings in period until very high radial order. Nonadiabatic solutions for these modes enable us to predict the relative visibility of them, and for  $l=1-5$ , the highest order modes seem to be more detectable. We find that the f-mode is pulsationally unstable for high  $l$  at the solar granulation space and time scales. The high helium needed to match the p-mode frequencies results in high central temperatures that give 13 SNU from the  ${}^8B$  and 2 SNU from the  ${}^7Be$  reactions.

**Table 1**  
**Solar Evolution Results**

Parameter	Without Diffusion	With Diffusion
Initial Y	0.300	0.298
Initial Z	0.020	0.020
Mixing Length Ratio	1.876	1.919
$T_{\text{eff}}$ (K)	5771.2	5770.4

**Table 2**  
**Solar Interior Results**

Parameter	Without Diffusion	With Diffusion
Central Y	0.6664	0.6672
Central Z	0.0206	0.0214
Central T ( $10^6\text{K}$ )	15.77	15.82
Central Density ( $g/cm^3$ )	165.4	165.6
$B^8$ Neutrinos (SNU)	12.6	13.0
$Be^7$ Neutrinos (SNU)	1.9	2.0

**Table 3**  
**Solar Envelope Results**

Parameter	Without Diffusion	With Diffusion
Surface Y	0.3003	0.2620
Surface Z	0.0200	0.0179
Convection Zone T ( $10^6\text{K}$ )	2.236	2.123
Convection Zone R ( $R_{\odot}$ )	0.717	0.721
Convection Zone M ( $M_{\odot}$ )	0.977	0.978




Figure 1. The hydrogen mass fraction in the mixture is plotted versus the internal mass fraction for the diffusion solar model.




Figure 2. Observed minus calculated p-mode frequencies are plotted versus the frequency for the no-diffusion model. These modes sample the model region below the convection zone.





Figure 3. Observed minus calculated p-mode frequencies are plotted versus the frequency for the no-diffusion model. These modes sample the convection zone.

Figure 4. Observed minus calculated p-mode frequencies are plotted versus the frequency for the diffusion model. These frequencies run a few microhertz above the similar plot for the no-diffusion model.




Figure 5. Observed minus calculated p-mode frequencies are plotted versus the frequency for the diffusion model. These modes sample the convection zone.

Figure 6. The  $\delta$  nonadiabatic frequency difference between the radial and quadrupole modes for radial orders 12 to 28 is plotted versus radial mode order for the diffusion model.




Figure 7. The period spacing between mode periods multiplied by  $\sqrt{l(l+1)}$  is plotted versus g-mode order. Fluctuations are due to our nonsmooth Brunt Väisälä frequency.

Figure 8. The work per zone over each pulsation cycle for the photospheric zones is plotted versus zone for the f mode with  $l=1400$ , near the solar granulation scale. The mode is pulsationally unstable because of the  $\kappa$  and  $\gamma$  effects. The work integral predicts an e-folding time to be about 7700 cycles, which, at the f mode period of 268 seconds, is only 25 days.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.